Neuroscience Letters, 172 (1994) 181–183
Elsevier Ireland Ltd.

AUTHOR INDEX

Abe, K., see Kobayashi, T., 147
Abraira, V., see Gaztelu, J.M., 31
Adamovich, S., Berkinblit, M., Smetanin, B., Fookson, O. and Poizner, H., Influence of movement speed on accuracy of pointing to memorized targets in 3D space, 171
Agid, Y., see Boka, G., 151
Alonso, R., see Welner, S.A., 89
Andersson, J., see Hoshi, Y., 129
Anglade, P., see Boka, G., 151
Annunziata, P., see Zhainazarov, A.B., 111
Ariano, M.A., see Larson, E.R., 101
Azzi, M., Nicot, A., Gully, D., Kitabgi, P., Bérod, A. and Rostène, W., Increase in neurotensin receptor expression in rat brain induced by chronic treatment with the nonpeptide neurotensin receptor antagonist SR 48692, 97

Barth, T., see Rex, A., 139
Bender, R., see Straube, A., 143
Benos, D.J., see Iwasa, K.H., 163
Bergström, M., see Hoshi, Y., 129
Berkinblit, M., see Adamovich, S., 171
Bérod, A., see Azzi, M., 97
Bischoff, A.-M., see Lalley, P.M., 59
Blanquet, P.R., see Torriglia, A., 125
Boka, G., Anglade, P., Wallach, D., Javoy-Agid, F., Agid, Y. and Hirsch, E.C., Immunocytochemical analysis of tumor necrosis factor and its receptors in Parkinson's disease, 151
Bolhuis, P.A., see van Gool, W.A., 122

Chen, S.T., Garey, L.J. and Jen, L.S., Bcl-2 proto-oncogene protein immunoreactivity in normally developing and axotomised rat retinas, 11
Chen, W.-T., Liu, W.-K. and Yen, S.-H., Expression of tau exon 8 in different species, 167
Cherubini, E., see Zhainazarov, A.B., 111
Ciofi, P., see Magoul, R., 134
Ciotti, M.T., see Volonte, C., 6
Clelland, C.A., see Lantos, P.L., 77
Coles, S.K., see Jodkowski, J.S., 67
Colling, S.B. and Wheal, H.V., Fast sodium action potentials are generated in the distal apical dendrites of rat hippocampal CA1 pyramidal cells, 73
Corda, M.G., see Giorgi, O., 35
Crunelli, V., see Juhász, G., 155
Cutrera, R.A., Ouarour, A. and Pévet, P., Effects of the 5-HT_{1a} receptor agonist 8-OH-DPAT and other non-photic stimuli on the circadian rhythm of wheel-running activity in hamsters under different constant conditions, 27

De Keyser, J., Wilczak, N. and Goossens, A., Insulin-like growth factor-I receptor densities in human frontal cortex and white matter during aging, in Alzheimer's disease, and in Huntington's disease, 93
Debarnot, F., see Rataud, J., 19
Deiana, A.M., see Giorgi, O., 35
Dick, T.E., see Jodkowski, J.S., 67
Dimaline, R., Evans, D., Forster, E. and Dockray, G.J., Stimulation of gastric somatostatin mRNA abundance by substance P in capsaicin-treated rats, 39
Dockray, G.J., see Dimaline, R., 39
Domeney, A.M., see Rex, A., 139

Emri, Z., see Juhász, G., 155
Endo, H., see Okamoto, M., 51
Evans, D., see Dimaline, R., 39

Fink, H., see Rex, A., 139
Fookson, O., see Adamovich, S., 171
Forster, E., see Dimaline, R., 39
Frotscher, M., see Straube, A., 143
Fuse, Y., see Kishida, H., 119

Garcia-Austt, E., see Gaztelu, J.M., 31
Garey, L.J., see Chen, S.T., 11
Gaztelu, J.M., Romero-Vives, M., Abraira, V. and Garcia-Austt, E., Hippocampal EEG theta power density is similar during slow-wave sleep and paradoxical sleep. A long-term study in rats, 31
Giorgi, O., Deiana, A.M., Salvadori, S., Lecca, D. and Corda, M.G., Developmental changes in the content of dopamine in the olfactory bulb of the European eel (*Anguilla anguilla*), 35
Goossens, A., see De Keyser, J., 93
Gully, D., see Azzi, M., 97

Higashida, H., see Kishida, H., 119
Hirsch, E.C., see Boka, G., 151
Hoshi, Y., Onoe, H., Watanabe, Y., Andersson, J., Bergström, M., Lilja, A., Långstöm, B. and Tamura, M., Non-synchronous behavior of neuronal activity, oxidative metabolism and blood supply during mental tasks in man, 129
Houpt, T.A., Philopena, J.M., Wessel, T.C., Joh, T.H. and Smith, G.P., Increased c-*fos* expression in nucleus of the solitary tract correlated with conditioned taste aversion to sucrose in rats, 1
Huang, C.-C., see Murphy, S., 42

Ichikawa, T., Light suppresses the activity of serotonin-immunoreactive neurons in the optic lobe of the swallowtail butterfly, 115
Ichimura, M., see Okamoto, M., 51
Iramina, K., see Yoshiura, T., 159
Iwasa, K.H., Mizuta, K., Lim, D.J., Benos, D.J. and Tachibana, M., Amiloride-sensitive channels in marginal cells in the stria vascularis of the guinea pig cochlea, 163
Iwata, M., see Parhar, I.S., 15

Javoy-Agid, F., see Boka, G., 151
Jeanny, J.C., see Torriglia, A., 125
Jen, L.S., see Chen, S.T., 11
Jodkowski, J.S., Coles, S.K. and Dick, T.E., A 'pneumotaxic centre' in rats, 67
Joh, T.H., see Houpt, T.A., 1
Johnson, J., see Lantos, P.L., 77
Juhász, G., Emri, Z., Kékesi, K.A., Salfay, O. and Crunelli, V., Blockade of thalamic $GABA_B$ receptors decreases EEG synchronization, 155

Kakei, S., Muto, N. and Shinoda, Y., Innervation of multiple neck

motor nuclei by single reticulospinal tract axons receiving tectal input in the upper cervical spinal cord, 85
Kékesi, K.A., see Juhász, G., 155
Kishida, H., Yamamoto, K., Fuse, Y., Noda, M. and Higashida, H., Activation of inward current associated with M-potassium current inhibition in m1-muscarinic receptor-transformed NG108-15 cells by KST-5452, a novel cognition enhancer, 119
Kitabgi, P., see Azzi, M., 97
Kobayashi, T., Abe, K., Saito, H. and Nishiyama, N., Effect of α_1-antichymotrypsin on the toxicity of β-amyloid fragment 25–40 in rat primary cultured neurons, 147
Koibuchi, N., see Parhar, I.S., 15

Långstöm, B., see Hoshi, Y., 129
Lajevardi, N., see Murphy, S., 42
Lalley, P.M., Bischoff, A.-M. and Richter, D.W., Serotonin 1A-receptor activation suppresses respiratory apneusis in the cat, 59
Lantos, P.L., Ovenstone, I.M.K., Johnson, J., Clelland, C.A., Roques, P. and Rossor, M.N., Lewy bodies in the brain of two members of a family with the 717 (Val to Ile) mutation of the amyloid precursor protein gene, 77
Laplante, P., see Welner, S.A., 89
Larson, E.R. and Ariano, M.A., Dopamine receptor binding on identified striatonigral neurons, 101
Lecca, D., see Giorgi, O., 35
Lilja, A., see Hoshi, Y., 129
Lim, D.J., see Iwasa, K.H., 163
Liu, W.-K., see Chen, W.-T., 167

Magoul, R., Ciofi, P. and Tramu, G., Visualization of an efferent projection route of the hypothalamic rat arcuate nucleus through the stria terminalis after labeling with carbocyanine dye (DiI) or proopiomelanocortin-immunohistochemistry, 134
Maragos, W.F. and Silverstein, F.S., Resistance to nitroprusside neurotoxicity in perinatal rat brain, 80
Mary, V., see Rataud, J., 19
Masuda, K., see Yoshiura, T., 159
Matsumoto, S. and Shimizu, T., Effects of isoprenaline on the responses of slowly adapting pulmonary stretch receptors to reduced lung compliance and to administered histamine, 47
Matsuo, S., see Shiraishi, Y., 24
Merlo, D., see Volonte, C., 6
Minamoto, Y., see Shibata, S., 107
Miura, M., see Takayama, K., 55
Mizuta, K., see Iwasa, K.H., 163
Mori, S., see Okamoto, M., 51
Murphy, S., Huang, C.-C., Lajevardi, N., Tammela, O., Wilson, D.F. and Pastuszko, A., Effect of hypoxia and reoxygenation on the activity of transglutaminase in brain of newborn piglets, 42
Muto, N., see Kakei, S., 85

Nakao, S., see Shiraishi, Y., 24
Nicot, A., see Azzi, M., 97
Nishiyama, N., see Kobayashi, T., 147
Nistri, A., see Zhainazarov, A.B., 111
Noda, M., see Kishida, H., 119

Okamoto, M., Mori, S., Ichimura, M. and Endo, H., Chondroitin sulfate proteoglycans protect cultured rat's cortical and hippocampal neurons from delayed cell death induced by excitatory amino acids, 51
Ono, M., see Shibata, S., 107
Onoe, H., see Hoshi, Y., 129
Otten, U., see Straube, A., 143
Ouarour, A., see Cutrera, R.A., 27
Ovenstone, I.M.K., see Lantos, P.L., 77

Parhar, I.S., Koibuchi, N., Sakai, M., Iwata, M. and Yamaoka, S., Gonadotropin-releasing hormone (GnRH): expression during salmon migration, 15
Pastuszko, A., see Murphy, S., 42
Peers, C., see Wyatt, C.N., 63
Pévet, P., see Cutrera, R.A., 27
Philopena, J.M., see Houpt, T.A., 1
Pierre, J.-A.S., see Welner, S.A., 89
Pinnock, R.D. and Woodruff, G.N., The non-peptide neurotensin receptor antagonist SR48962 is not a potent antagonist of neurotensin(8–13) responses of rat substantia nigra neurones in vitro, 175
Poizner, H., see Adamovich, S., 171
Pratt, J., see Rataud, J., 19

Rataud, J., Debarnot, F., Mary, V., Pratt, J. and Stutzmann, J.-M., Comparative study of voltage-sensitive sodium channel blockers in focal ischaemia and electric convulsions in rodents, 19
Rex, A., Barth, T., Voigt, J.-P., Domeney, A.M. and Fink, H., Effects of cholecystokinin tetrapeptide and sulfated cholecystokinin octapeptide in rat models of anxiety, 139
Richter, D.W., see Lalley, P.M., 59
Romero-Vives, M., see Gaztelu, J.M., 31
Roques, P., see Lantos, P.L., 77
Rossor, M.N., see Lantos, P.L., 77
Rostène, W., see Azzi, M., 97

Saito, H., see Kobayashi, T., 147
Sakai, M., see Parhar, I.S., 15
Salfay, O., see Juhász, G., 155
Salvadori, S., see Giorgi, O., 35
Schenk, D.B., see van Gool, W.A., 122
Shibata, S., Minamoto, Y., Ono, M. and Watanabe, S., Aging impairs methamphetamine-induced free-running and anticipatory locomotor activity rhythms in rats, 107
Shimizu, T., see Matsumoto, S., 47
Shinoda, Y., see Kakei, S., 85
Shiraishi, Y., Matsuo, S. and Nakao, S., Medium-lead burst neurons related to vertical saccades in cat Forel's field H: input from vestibular nucleus, 24
Silverstein, F.S., see Maragos, W.F., 80
Smetanin, B., see Adamovich, S., 171
Smith, G.P., see Houpt, T.A., 1
Straube, A., Bender, R., Otten, U. and Frotscher, M., Expression of the cholecystokinin gene in rat hippocampal interneurons is independent of extrinsic input, 143
Stutzmann, J.-M., see Rataud, J., 19

Tachibana, M., see Iwasa, K.H., 163
Takayama, K., Xiong, Y. and Miura, M., Neuronal expression of Fos protein in the paraventricular nucleus of the hypothalamus after i.p. injection of ulcergenic cinchophen, 55
Tammela, O., see Murphy, S., 42
Tamura, M., see Hoshi, Y., 129
Toneatto, S., see Zhainazarov, A.B., 111
Torriglia, A., Jeanny, J.C. and Blanquet, P.R., Immunohistochemical analysis of fibroblast growth factor receptor in bovine retina, 125
Tramu, G., see Magoul, R., 134

Ueno, S., see Yoshiura, T., 159

van Gool, W.A., Schenk, D.B. and Bolhuis, P.A., Concentrations of amyloid-β protein in cerebrospinal fluid increase with age in patients free from neurodegenerative disease, 122
Voigt, J.-P., see Rex, A., 139
Volonte, C., Ciotti, M.T. and Merlo, D., LiCl promotes survival of GABAergic neurons from cerebellum and cerebral cortex: LiCl induces survival of GABAergic neurons, 6

Wallach, D., see Boka, G., 151
Watanabe, S., see Shibata, S., 107
Watanabe, Y., see Hoshi, Y., 129
Weir, E.K., see Wyatt, C.N., 63
Welner, S.A., Laplante, P., Pierre, J.-A.S. and Alonso, R., Excitatory and inhibitory amino acids in the cerebral cortex of nucleus basalis magnocellularis lesioned rats, 89
Wessel, T.C., see Houpt, T.A., 1
Wheal, H.V., see Colling, S.B., 73
Wilczak, N., see De Keyser, J., 93
Wilson, D.F., see Murphy, S., 42
Woodruff, G.N., see Pinnock, R.D., 175
Wyatt, C.N., Weir, E.K. and Peers, C., Diphenylene iodonium blocks K^+ and Ca^{2+} currents in type I cells isolated from the neonatal rat carotid body, 63

Xiong, Y., see Takayama, K., 55

Yamamoto, K., see Kishida, H., 119
Yamaoka, S., see Parhar, I.S., 15
Yen, S.-H., see Chen, W.-T., 167
Yoshiura, T., Ueno, S., Iramina, K. and Masuda, K., Effects of stimulation side on human middle latency auditory evoked magnetic fields, 159

Zhainazarov, A.B., Annunziata, P., Toneatto, S., Cherubini, E. and Nistri, A., Serum fractions from amyotrophic lateral sclerosis patients depress voltage-activated Ca^{2+} currents of rat cerebellar granule cells in culture, 111